

Bosonization Theory of Excitons in One-Dimensional Narrow Gap Semiconductors

H. C. Lee¹ and S. -R. Eric Yang²

BK21 Physics Research Division and Institute of Basic Science, Sung Kyun Kwan University, Suwon, 440-746 Korea¹
and

Department of Physics, Korea University, Seoul, Korea²

Excitons in one-dimensional narrow gap semiconductors of anti-crossing quantum Hall edge states are investigated using a bosonization method. The excitonic states are studied by mapping the problem into a non-integrable sine-Gordon type model. We also find that many-body interactions lead to a strong enhancement of the band gap. We have estimated when an exciton instability may occur.

PACS numbers: 71.27.+a, 71.35.-y, 71.35.Cc

One-dimensional (1D) narrow gap semiconductors can be realized in anti-crossing quantum Hall edge states¹ and nanotubes². The gaps in these systems are *single* particle gaps and *not* many body gaps. Theoretically, they provide the unusual condition that the bare band gap t is much *smaller* than the characteristic Coulomb energy scale E_c . Moreover, strong quantum fluctuations are present in these systems, reflecting the 1D character. The ground state of these systems may be unstable against the spontaneous formation of excitons if the exciton binding energy exceeds the band gap^{3,4}. In three dimensional semiconductors excitons can be treated successfully by solving the Bethe-Salpeter equation⁴. However, in the strong coupling regime of 1D systems, the perturbative approaches are not expected to be reliable due to large quantum fluctuations. If the Coulomb scale is much larger than the gap one might naively expect that exciton instability would occur. However, this simple picture neglects screening which is expected to be large due to the smallness of the gap. It is unclear whether a bound state of an "electron" and a "hole" can exist in the presence of strong quantum fluctuations. Not much is known about the physics of excitons in 1D narrow gap semiconductors.

This problem can be addressed within a bosonization approach, which is applicable as long as the characteristic energy scale of the problem is smaller than the band width W . Thus bosonization provides a natural framework for studying 1D excitons. In this work we will consider anti-crossing of quantum Hall edge states in the barrier region of between two 2D electron gases. Assume that the applied magnetic field is sufficiently strong that the system is spin-polarized with the filling factor $\nu = 1$ and that the Fermi level is in the gap of anti-crossing edge states. We find that the problem can be mapped into a sine-Gordon (sG) model with an extra term representing the long range Coulomb interaction. Due to this extra term the model becomes *non-integrable* and an exact solution is unavailable. However, the non-linear cosine term of this sG model can be expanded provided that $\delta_{ex} \sim \frac{v/v_0}{\ln W/t} < 1$, where v is the Fermi velocity, $v_0 = e^2/\epsilon\hbar$, $W = \hbar v/a$, and a is the smallest length

scale in the problem⁶. In this regime a perturbative theory may be applied to calculate excitonic energies. Our work also indicates that a large enhancement of band gap occurs. Our perturbative approach suggests an approximate estimate for when an exciton instability may occur: $\delta_{ex} > 1$. (We cannot exclude the possibility that the higher order corrections neglected in our perturbative calculation may work against such an instability).

The system is modeled by the following Hamiltonian⁵

$$\begin{aligned} H &= H_0 + H_{\text{coul}} + H_t, \\ H_0 &= v \int dx \left[-i\psi_R^\dagger \partial_x \psi_R + i\psi_L^\dagger \partial_x \psi_L \right] \\ &= \pi v \int dx \left[\rho_R^2 + \rho_L^2 \right] \end{aligned} \quad (1)$$

$$\begin{aligned} H_{\text{coul}} &= \int dx dy \left[\frac{V(x-y)}{2} (\rho_R(x)\rho_R(y) + \rho_L(x)\rho_L(y)) \right. \\ &\quad \left. + V(x-y)(\rho_L(x)\rho_R(y)) \right] \end{aligned}$$

$$H_t = -t \int dx \left[\psi_R^\dagger(x)\psi_L(x) + \text{H.C.} \right], \quad (2)$$

The operators $\psi_R(\psi_L)$ is the right-moving (left-moving) edge *electron* operators. $\rho_R =: \psi_R^\dagger \psi_R$: is the (normal-ordered) right-moving edge electron density operator (ρ_L is similarly defined). In practice, we can take $V(x) = \frac{e^2}{\epsilon} \frac{1}{\sqrt{x^2+a^2}}$ and the Coulomb matrix element is $V(k) = \frac{2e^2}{\epsilon} K_0(a|k|)$. The tunneling between the right-moving and left-moving is modeled by H_t . The tunneling amplitude t is assumed to be significant only near the Fermi wave vector k_F (The Fermi wavevector is set to zero from now on). The cut-off value of the wave vector is $1/a$. Note that a single particle gap opens up near the Fermi points due to this tunneling term. The commutation relations of density operators in momentum space are given by⁵ $[\rho_R(q), \rho_R(q')] = -\frac{q}{2\pi} \delta_{q+q'}$, $[\rho_L(q), \rho_L(q')] = +\frac{q}{2\pi} \delta_{q+q'}$. The interacting electron systems can be bosonized in a standard way. The explicit relations between the electron operators and the bosonic variable ϕ_R, ϕ_L are $\rho_R = \frac{1}{2\pi} \partial_x \phi_R$, $\rho_L = \frac{1}{2\pi} \partial_x \phi_L$, $\psi_R = \frac{1}{\sqrt{2\pi a}} e^{i\phi_R}$, $\psi_L = \frac{1}{\sqrt{2\pi a}} e^{-i\phi_L}$. It is conve-

nient to introduce the conjugate fields $\phi_{\pm}$: $\phi_{\pm} \equiv \phi_R \pm \phi_L$. The effective bosonized action in imaginary time reads

$$S = \int dx d\tau \left[\frac{v}{8\pi} ((\partial_x \phi_+)^2 + (\partial_x \phi_-)^2) + \frac{i}{4\pi} \partial_\tau \phi_+ \partial_x \phi_- \right] + \frac{1}{8\pi^2} \int dx dy d\tau V(x-y) \partial_x \phi_+(x) \partial_y \phi_+(y) - \frac{t}{2\pi a} \int dx d\tau \left[e^{i\phi_+(x)} + \text{H.C.} \right]. \quad (3)$$

The ϕ_- can be integrated out, leaving us with

$$S = \frac{1}{8\pi} \int \frac{dk d\omega}{(2\pi)^2} \left[vk^2 \left(1 + \frac{V(k)}{\pi v} \right) + \frac{\omega^2}{v} \right] \times \phi_+(\omega, k) \phi_+(-\omega, -k) - \frac{t}{\pi a} \int dx d\tau \cos(\phi_+(x, \tau)). \quad (4)$$

The above action looks very similar to the sG model⁷, except for the momentum-dependent Coulomb interaction $V(k)$. If $V(k)$ were momentum independent (local interaction in real space), the action would be exactly that of sG model. The Euclidean action of the sine-Gordon model is given by⁷⁻⁹

$$A_{SG} = \int d^2x \left[\frac{1}{16\pi} (\partial_\mu \phi)^2 - 2\mu \cos(\beta\phi) \right], \quad (5)$$

where the speed of "light" has been set to unity¹⁰. It is useful to show the equivalence between the above sG action and our action (4) for the short range interaction ($V(k) = V$ is constant). This is achieved by the relations

$$\phi = \phi_+ [4(1 + \frac{V}{\pi v})]^{1/4}, \quad \beta = [4(1 + \frac{V}{\pi v})]^{-1/4}, \quad \mu = \frac{t}{\pi a v (1 + V/\pi v)^{1/2}}. \quad (6)$$

From the above relations it is clear that the *strong* coupling regime of the original electron system (large V) is mapped to the *weak* coupling regime (small β) of sG model.

Before we investigate the long-range case it is instructive to review some known physical properties of sG model, which is exactly solvable. The spectrum of sG model consist of the breathers (B_n), the soliton, and the anti-soliton. The breathers are the bound states of the soliton and anti-soliton⁷⁻⁹. The lightest bound state B_1 coincides with the fundamental boson ϕ of the sG model (5), and it is associated with the perturbative treatment. The number of breather is given by $n = 1, 2, 3, \dots < 1/\xi$, where $\xi = \frac{\beta^2}{1-\beta^2}$. Note that even for infinitesimally small short range repulsion V , at least one breather exists. The breather energy (excitation energy) is given by

$$m = 2M \sin \frac{\pi\xi}{2}, \quad (7)$$

where M is the energy of soliton. The parameter μ and the soliton energy M are related through

$$\mu = \frac{\Gamma(\beta^2)}{\pi\Gamma(1-\beta^2)} \left[M \frac{\sqrt{\pi}\Gamma(\frac{1+\xi}{2})}{2\Gamma(\frac{\xi}{2})} \right]^{2-2\beta^2}. \quad (8)$$

Note that in the $\beta \rightarrow 0$ limit, $M \sim v_\rho^2 \mu / \beta^2 \rightarrow \infty$ and $m \sim v_\rho^2 (\mu / \beta^2) \beta^4 \sim v_\rho^2 \mu \beta^2 \sim t/va$, where "the speed of light" v_ρ has been reinstated for clarity¹⁰. In other words, in the large V limit the soliton becomes very massive and leads to a large enhancement of the band gap. In contrast, the lightest breather energy approaches a constant value given by the coefficient of the cosine term of Eq.(4).

In the *absence* of the Coulomb interaction the exact value of the single particle gap should be t . However, the bosonized action (4) contain the factor a , and it is unclear how this factor disappears in the final result for the gap. Let us try to understand how this happens. The crucial fact is that the cosine term of the sG action (5) is normal ordered and it gives the dimension to the cosine operator. The exact results Eq.(7, 8)^{8,9} were obtained with

$$\mu^2 \langle : \cos(\beta\phi(x)) :: \cos(\beta\phi(y)) : \rangle \rightarrow \frac{1}{2} \frac{\mu^2}{|x-y|^{4\beta^2}}, \quad (9)$$

as $|x-y| \rightarrow 0$. In the absence of Coulomb interaction, corresponding to $\beta^2 = 1/2$, the same correlation functions (two-point function of tunneling term) can be computed exactly when expressed in terms of electron operators

$$\langle [t\psi_R^\dagger(x)\psi_L(x)][t\psi_L^\dagger(y)\psi_R(y)] \rangle = \frac{t^2}{|x-y|^2}. \quad (10)$$

The formal application of bosonization formula to this electron tunneling term gives $\frac{t}{\pi a} \cos(\beta\phi)$ as in our action (4). As it stands it is not normal ordered, and consequently the short-distance singularity must be regularized in the calculation of correlation function.

$$\left(\frac{t}{2\pi a} \right)^2 \langle e^{i\beta\phi(x)} e^{-i\beta\phi(y)} \rangle = \left(\frac{t}{2\pi a} \right)^2 \exp \left[-\beta^2 (\langle \phi(0)\phi(0) \rangle - \langle \phi(x)\phi(y) \rangle) \right]. \quad (11)$$

We have to regularize $\langle \phi(0)\phi(0) \rangle$ as $\langle \phi(0)\phi(a) \rangle = 2 \ln \frac{L^2}{a^2}$, where L is the system size. L always appears in the definition of propagator but they are canceled for the physical correlation functions (see below). Since $\langle \phi(x)\phi(y) \rangle = 2 \ln \frac{L^2}{|x-y|^2}$ we obtain

$$\left(\frac{t}{2\pi a} \right)^2 \langle e^{i\beta\phi(x)} e^{-i\beta\phi(y)} \rangle = \left(\frac{t}{2\pi a} \right)^2 \frac{1}{(|x-y|/a)^{4\beta^2}}. \quad (12)$$

Only at $\beta^2 = 1/2$, the length scale a drops out. Comparing Eq.(9) with Eq.(12), the t can identified with μ .

Then by applying formula Eq.(8) at $\beta^2 = 1/2$, we find that $t = M$, which is the expected result. The above argument checks the consistency of the bosonization formulation of our problem. We emphasize again, the above results holds only if the Coulomb interaction is absent.

When the long range Coulomb interaction is present the model becomes *non-integrable* and an exact solution is unavailable. But, under certain conditions, the cosine term of the action (4) can be expanded and excitons may be studied perturbatively. (See the perturbative calculation below). Expanding the cosine term of (4) up to the 4th order, we get an effective action of exciton states.

$$S \approx \frac{1}{8\pi} \int \frac{dkd\omega}{(2\pi)^2} \left[vk^2 \left(1 + \frac{V(k)}{\pi v} \right) + \frac{\omega^2}{v} + \frac{4t}{\hbar a} \right] \times \phi_+(\omega, k) \phi_+(-\omega, -k) - \frac{1}{4!} \frac{t}{\pi \hbar a} \int dx d\tau : \phi_+^4 : . \quad (13)$$

At this point, we note that the parameter $\frac{t}{\pi a}$ in action (4) should be understood as a renormalized quantity. The application of bosonization formula to the fermion bilinear mass term assumes the implicit normal ordering¹¹. Such a normal ordering sums the tadpole diagrams¹² (see Fig.1) and the effect of the normal ordering is contained in the cut-off parameter $\frac{1}{2\pi a}$. Therefore, in the perturbative expansion of cosine term we have to exclude all the tadpole type diagrams, and then all other terms of perturbative expansion are finite and well-defined. In analogy with the sG model, we will associate the soliton-like (quantum) solution of (4) with the renormalized electrons, and the breather-like ϕ_+ mode with the excitonic electron-hole bound state.

If the quantum correction due to ϕ_+^4 is neglected, the zeroth order dispersion relation of exciton state can be read off from the quadratic part of (13).

$$E^{(0)}(k) = \hbar \left[v^2 k^2 \left(1 + \frac{v_0}{2v} \ln \frac{1}{a|k|} \right) + \frac{4tv}{\hbar a} \right]^{1/2}. \quad (14)$$

The energy gap is given by

$$\Delta_{ex}^{(0)} = \sqrt{\frac{4\hbar tv}{a}} = 2\sqrt{tW}. \quad (15)$$

Next, we compute the correction to the exciton excitation energy due to the quantum fluctuation of ϕ_+^4 term. The first such correction appears in the second order perturbation of ϕ_+^4 . Because the ϕ_+^4 term is normal ordered there are no tadpole diagrams. Only the "sunset" diagram (see Fig.1) contributes, which is both ultraviolet and infrared convergent.




FIG. 1. (a) is one of the diagrams contributing to the normal ordering, and (b) is the "sunset" diagram. The solid lines are the boson propagator $\langle \phi_+ \phi_+ \rangle$.

It contributes a *negative* correction to the gap as can be checked by a simple calculation $e^{-(\Delta_{ex}^{(0)})^2 \phi^2 (1 + \lambda^2 \phi^2 \langle \phi^3 \phi^3 \rangle)} \sim e^{-((\Delta_{ex}^{(0)})^2 - \lambda^2 \langle \phi^3 \phi^3 \rangle) \phi^2}$, where $\lambda \sim t/\pi \hbar a$. The symmetry factor of the sun-set diagram is $\frac{1}{4!} \frac{1}{4!} 4 \cdot 4 \cdot 3 \cdot 2 = 1/6$. In terms of self-energy the sunset diagram gives

$$\Sigma(i\omega, k) = \frac{1}{6} \left(\frac{t}{\pi \hbar a} \right)^2 \int dx d\tau [D(x, \tau)]^3 e^{ikx + i\omega \tau},$$

$$D(x, \tau) = \int \frac{d\omega dk}{(2\pi)^2} \frac{v e^{ikx + i\omega \tau}}{\omega^2 + v^2 k^2 \left(1 + \frac{\alpha}{2} \ln \frac{1}{|k|a} \right) + \frac{4tv}{\hbar a}}. \quad (16)$$

The excitation energy which includes the above quantum correction is given by

$$\Delta_{ex} = \left[\frac{4\hbar tv}{a} - 8\pi \hbar^2 v \Sigma(0, 0) \right]^{1/2}. \quad (17)$$

Write the above equation in the form

$$\Delta_{ex}^2 = (\Delta_{ex}^{(0)})^2 \left[1 - \frac{\Sigma(0, 0)}{t/(2\pi \hbar a)} \right]. \quad (18)$$

The second term in the bracket of Eq.(18) is the quantum correction in the dimensionless form. Explicitly, (y_1, y_2, η_1, η_2 are dimensionless).

$$\delta_{ex} = \frac{\Sigma(0, 0)}{t/(2\pi \hbar a)} = \frac{1}{4\pi} \int dy_1 dy_2 [D(y_1, y_2)]^3,$$

$$D(y_1, y_2) = \int \frac{d\eta_1 d\eta_2}{(2\pi)^2} \frac{e^{i\eta_1 y_1 + i\eta_2 y_2}}{4 + \eta_1^2 + \eta_2^2 f(|\eta_2|)}, \quad (19)$$

where $f(|\eta_2|) = 1 + \frac{v_0}{2v} \ln \left[\frac{1}{|\eta_2|} \sqrt{\frac{W}{t}} \right]$. We may extract the effective expansion parameter by rescaling. It is easy to see that

$$\delta_{ex} \sim \frac{\delta \Sigma(0, 0)}{t/(2\pi \hbar a)} \sim \frac{v/v_0}{\ln W/t}. \quad (20)$$

We note that, in the perturbative regime $\delta_{ex} \rightarrow 0$ the excitation energy approaches the value of the coefficient of the cosine term. This same feature also appeared in the exactly solvable sine-Gordon model. Our perturbative treatment for the excitation states is thus valid when $\delta_{ex} < 1$. This result suggests the possibility that when $\delta_{ex} > 1$ is satisfied exciton instability may occur. However, we cannot exclude the possibility that the higher

order corrections neglected in our perturbative calculation may prohibit such an instability.

The action Eq.(4) supports topological soliton excitations due to the topologically inequivalent vacuum of the cosine potential⁷. Thus, we can *not* use the perturbative expansion of the cosine term in the study of the soliton excitation. The original sine-Gordon model can be solved exactly, and the energy gaps of soliton and breather are known exactly (see Eq.(7), Eq.(8))^{7,9}. Our action (4) is not exactly solvable, and here we will give only an estimate of the gap of soliton excitation. In estimating the gap of soliton excitation, it is sufficient to consider the static soliton. In our perturbative regime we may neglect 1 compared to $\frac{V(k)}{\pi v}$. Then, in real space, the energy functional for static solution reads

$$E[\phi_+(x)] = \frac{1}{8\pi} \int dx dy \frac{e^2}{\epsilon|x-y|} \partial_x \phi_+(x) \partial_y \phi_+(y) + \frac{t}{\pi a} \int dx (1 - \cos(\phi_+(x))), \quad (21)$$

(A constant term is added to make the expression positive definite). We define a dimensionless variable $\bar{x}$, $\bar{x} = x \left(\frac{t}{e^2 \epsilon a}\right)^{1/2} = x \left(\frac{t}{E_c}\right)^{1/2} \frac{1}{a}$. Then, the energy functional becomes

$$E[\phi_+(x)] = \sqrt{\frac{e^2 t}{\epsilon a}} \left[\frac{1}{8\pi} \int d\bar{x} d\bar{y} \frac{1}{|\bar{x} - \bar{y}|} \partial_{\bar{x}} \phi_+(\bar{x}) \partial_{\bar{y}} \phi_+(\bar{y}) + \frac{1}{\pi} \int d\bar{x} (1 - \cos(\phi_+(\bar{x}))) \right]. \quad (22)$$

The coefficient in front of bracket has the dimension of energy and it gives a characteristic gap scale of solitons. The ground state, which sits at a minimum of the cosine potential, has zero classical energy. The soliton excitation connects two adjacent classical ground states, and according the above estimate the gap of soliton E_G is of the order $\sqrt{\frac{e^2 t}{\epsilon a}} = \sqrt{E_c t}$. Since we expect $E_c \gg t$ the Coulomb interaction enhances the value of band gap significantly.

In summary, we have studied excitons in 1D narrow gap semiconductors of two anticrossing quantum Hall edges. According to our perturbative approach the exciton state may lie in the gap when $\delta_{ex} < 1$. Our study indicates that many body interactions enhances the value of band gap significantly. Our result suggests that an exciton instability may occur when $\delta_{ex} \sim 1$. However, it

is desirable to calculate how the higher order corrections neglected in our perturbative calculation may change this condition. The actual values of v , and t in anticrossing edges states are not known well and it is difficult to estimate precisely the actual value of δ_{ex} ¹³. Experimental observation of excitons and the investigation of exciton instability would be most interesting.

We are grateful to Al. B. Zamolodchikov and M.P.A. Fisher for useful comments. H. C. L. was supported by the Korea Science and Engineering Foundation (KOSEF) through the grant No. 1999-2-11400-005-5, and by the Ministry of Education through Brain Korea 21 SNU-SKKU Program. S.R.E.Y was supported by Korea Research Foundation Grant KRF-2000- 015-DP0125.

-
- ¹ W. Kang, H. L. Stomer, K. W. Baldwin, L. N. Pfeiffer, and K. W. West, *Nature*, **403**, 59 (2000).
 - ² T. W. Ebbesen, *Physics Today*, **49** (6), 26(1996); N. Hamada, S. Sawada, and A. Oshiyama, *Phys. Rev. Lett* **68**, 1579 (1992).
 - ³ B. I. Halperin and T. M. Rice, *Rev. Mod. Phys.* **40**, 755 (1968).
 - ⁴ L. J. Sham and T.V. Ramakrishnan, in *Physics of Semimetals and Narrow gap Semiconductors*, edited by D.L. Carter and R.T. Bate (Pergamon, New York, 1971), p.219.
 - ⁵ X. G. Wen, *Phys. Rev. B* **41**, 12838 (1990); *Phys. Rev. B*, **44**, 5708 (1991); *Int. Journ. Mod. Phys. B*, **6**, (1992).
 - ⁶ There are several small lengths in this problem: the transverse dimension of the quasi-1D system w , the magnetic length ℓ , and the width of the barrier a . They are all of order 100 Å. In this paper they will all be denoted by a .
 - ⁷ R. Rajaraman, *Solitons and Instantons* (North-Holland, New York, 1987).
 - ⁸ S. Lukyanov and A. Zamolodchikov, *Nucl. Phys. B* **493**, 571 (1997).
 - ⁹ Al. B. Zamolodchikov, *Int. Journ. Mod. Phys. B* **A 10**, 1125 (1995).
 - ¹⁰ The explicit expression of the "speed of light" is $v_\rho = v(1 + V/\pi v)^{1/2}$. Note its dependence on V .
 - ¹¹ S. Mandelstam, *Phys. Rev. D* **11**, 3026 (1975).
 - ¹² S. Coleman, *Phys. Rev. D* **11**, 2088 (1975).
 - ¹³ In our estimate it is likely that $0.1 < \delta_{ex} < 1$